

ONLINE FIRST | LESS IS MORE

The “Top 5” Lists in Primary Care

Meeting the Responsibility of Professionalism

The Good Stewardship Working Group

Background: Physicians can adhere to the principles of professionalism by practicing high-quality, evidence-based care and advocating for just and cost-effective distribution of finite clinical resources. To promote these principles, the National Physicians Alliance (NPA) initiated a project titled “Promoting Good Stewardship in Clinical Practice” that aimed to develop a list of the top 5 activities in family medicine, internal medicine, and pediatrics where the quality of care could be improved.

Methods: Working groups of NPA members in each of the 3 primary care specialties agreed that an ideal activity would be one that was common in primary care practice, that was strongly supported by the evidence, and that would lead to significant health benefits and reduce risks, harms, and costs. A modification of nominal group process was used to generate a preliminary list of activities. A first round of field testing was conducted with 83 primary care physicians, and a second round of field testing with an additional 172 physicians.

Results: The first round of field testing resulted in 1 activity being deleted from the family medicine list. Sup-

port for the remaining activities was strong. The second round of field testing showed strong support for all activities. The family medicine and internal medicine groups independently selected 3 activities that were the same, so the final lists reflect 12 unique activities that could improve clinical care.

Conclusions: Physician panels in the primary care specialties of family medicine, internal medicine, and pediatrics identified common clinical activities that could lead to higher quality care and better use of finite clinical resources. Field testing showed support among physicians for the evidence supporting the activities, the potential positive impact on medical care quality and cost, and the ease with which the activities could be performed. We recommend that these “Top 5” lists of activities be implemented in primary care practice across the United States.

Arch Intern Med.

Published online May 23, 2011.

doi:10.1001/archinternmed.2011.231

IN 2009, THE AMERICAN BOARD OF Internal Medicine Foundation launched “Putting the Charter into Practice,” a program providing small grants to advance principles of professional commitment in medicine, in keeping with the Physician Charter,¹ a document jointly issued in 2002 by the American Board of Internal Medicine Foundation, the American College of Physicians Foundation, and the European Federation of Internal Medicine. These principles include improving patients’ access to high-quality care, practicing evidence-based care, advocating for just and cost-effective distribution of finite resources, and maintaining trust by minimizing conflicts of interest.

The National Physicians Alliance (NPA) was founded in 2005 and today represents 22 000 members across specialties and across the United States. The NPA’s primary mission is to ensure affordable,

high-quality health care for all; the organization is committed to building a sustained network of physician leaders dedicated to achieving this goal.

The NPA was awarded a grant by the American Board of Internal Medicine Foundation to develop and disseminate 5-activity lists of evidence-based, quality-improving, resource-sparing activities that could be incorporated into the practices of primary care providers in family medicine, internal medicine, and pediatrics. Each activity was to be well supported by evidence, have beneficial effects on patient health by improving treatment and/or reducing risks, and, where possible, reduce costs of care.

Simultaneously, an editorial on the medical ethics of health care reform by Howard Brody² called on each medical specialty to generate top-5 lists of diagnostic tests or treatments that are commonly ordered but that offer limited benefits or

Author Affiliations: All members of The Good Stewardship Working Group were authors. Their affiliations are on page E5.

Group Information: The Good Stewardship Working Group members are listed at the end of this article.

carry risks that outweigh their benefits. Brody asserted that by focusing on the top 5 examples of the most egregious causes of waste in each specialty, the medical profession could demonstrate to a skeptical and concerned public that high-quality care and efficient use of resources are complementary.

To compile an evidence-based top-5 list that would improve quality of care in the context of limited resources, the NPA initiated a project entitled “Promoting Good Stewardship in Clinical Practice.”

METHODS

The project began by assembling working groups in each of the 3 primary care specialties. An e-mail solicitation was sent to all NPA members, inviting participation by interested primary care physicians. The project director (S.R.S.) selected a chairperson and 4 other physicians for each working group from among the respondents to the e-mail solicitation. Selection took into account geographical, gender, and racial and/or ethnic diversity and yielded 15 members: 10 women, 2 Asian Americans, 1 African American, 1 Hispanic American, and 2 physicians from rural areas.

Each working group met in a series of teleconference calls to delineate the candidate top-5 activities. The groups agreed that an ideal activity would be one that was common in primary care practice, that was strongly supported by the evidence, and that would lead to significant health benefits, reduce risks and harms to patients and communities, and reduce costs.

A modification of nominal group process was used to generate an initial list of top-5 activities.³ A shared Google document was used to generate the list and conduct the voting online in real time.

A research assistant reviewed the literature to find evidence supporting or refuting the candidate top-5 activities and reported this to the working groups. Working groups modified their candidate activities in response to these findings, and each group generated a preliminary list of top-5 activities for field testing.

Working group members were asked to recruit 3 to 5 physicians from their specialties to serve as initial field testers. Project goals were explained to the 83 initial testers, and each tester consented to participate. Each completed an online survey in which they rated candidate activities for their specialty on 5 parameters: (1) frequency with which they engaged in this activity in their practice; (2) the potential impact of the activity on quality of care; (3) the potential impact of the activity on cost of care; (4) the strength of the evidence supporting the activity; and (5) the ease or difficulty of implementing the activity in their own practice.

Physicians in the 3 primary care specialties were recruited for the second round of field testing through a general e-mail invitation sent to all NPA members. The 172 second-round testers completed the same survey as the initial testers.

The activity was considered well supported if, on average, a majority of the survey respondents rated the activity in the 2 highest categories on the 5-point scale (ie, “significant” or “large” on the quality-of-care and economic-impact parameters, “somewhat strong” or “very strong” on the evidence parameter, and “not difficult” or “easy” on the ease-of-implementation parameter).

The first question on the survey asked the respondents “How often are you confronted with patients where you would have to decide whether or not to do the following tests or procedures?” The responses, however, seemed to indicate that a substantial number of the respondents misinterpreted the question and reported how often they *followed* the recommendation.

The question was rephrased in the beta test to “How often do you need to make a decision whether or not to order the following tests, procedures, or treatments?” The problem with misinterpretation persisted. Therefore, the data from the first question on the survey, titled “Frequency Encountered,” were deemed to be invalid and were not considered further in deciding whether to keep the item.

Members of the working groups were unanimous in their belief that most of the candidate activities were sufficiently common in primary care practice to be included. For clinical encounters that were not as common, such as children presenting with minor head trauma, the working group believed that this was important enough to include anyway.

RESULTS

In the first round of testing, the 83 field testers generally rated the items favorably (**Table**), but 1 activity in the family medicine list—not doing routine blood chemical analysis for asymptomatic, healthy adults—showed weak support. While 70.3% felt that the evidence supporting the recommendation was somewhat strong or very strong, only 37.0% felt that adhering to the recommendation was quite important or critically important to the quality of care, and only 48.1% felt that it would be easy or not difficult to implement. Therefore, the activity was dropped, and another candidate activity (not routinely prescribing antibiotics for mild to moderate sinusitis) replaced it for the second field test. All other activities underwent only minor wording changes, based on the feedback of the first field testers.

A second round of field testing was performed on the revised lists with a new group of 172 primary care physicians. The results of the second field test are also listed in **Table**. These testers endorsed the activities on the revised list.

The final top-5 lists are presented in the **Figures 1, 2, and 3**. Although working groups functioned independently of one another, some activities arose in more than 1 group. The family medicine and internal medicine groups independently selected 3 activities that were the same, so the final lists reflect 12 unique activities. This commonality across specialties reinforced the importance and relevance of addressing overuse of these activities. When substantively similar items arose in different groups, one working group was designated as the lead on that item, modified the wording in response to field testers from all relevant specialties, and returned the activity to the second working group for ratification.

COMMENT

Physician panels in the primary care specialties of family medicine, internal medicine, and pediatrics identified common clinical activities where changes in practice could lead to higher-quality care and better use of finite clinical resources. Field testing showed support among physicians for the evidence supporting the recommendations, the potential positive impact on quality and cost, and the ease with which the recommendations could be implemented.

Table. Field Testers Who Agreed or Strongly Agreed on the Importance of the Clinical Activity

Practice Activity	Testing Group	Field Testers Who Agreed or Strongly Agreed on Importance, %				
		Frequency Encountered	Quality of Care	Economic Impact	Strength of Evidence	Ease of Implementation
Family Medicine						
Imaging for low back pain	Alpha	55.5	51.8	85.1	100.0	70.3
	Beta	38.2	41.9	81.8	96.4	83.6
Routine laboratory studies	Alpha	74.1	37.0	59.2	70.3	48.1
Antibiotics for sinusitis	Beta	38.2	63.6	60.0	94.6	70.9
Annual ECGs	Alpha	44.4	40.7	48.1	92.6	88.9
	Beta	29.0	40.0	54.5	92.7	96.3
Pap test for patients <21 y	Alpha	40.7	74.1	59.2	96.3	74.0
	Beta	36.4	51.0	50.9	96.3	89.1
DEXA scans younger patients	Alpha	40.7	55.5	66.7	85.1	81.4
	Beta	25.4	32.7	60.0	90.9	83.7
Internal Medicine						
Imaging for low back pain	Alpha	65.5	75.8	82.7	100.0	75.8
	Beta	40.0	49.0	74.0	90.0	80.0
Routine laboratory studies	Alpha	72.1	51.7	62.1	68.9	44.8
	Beta	54.0	40.0	54.0	78.0	60.0
Annual ECGs	Alpha	58.6	69.0	62.0	89.7	89.6
	Beta	26.0	40.0	42.0	84.0	88.0
Generic statins	Alpha	75.9	79.3	86.2	89.7	93.1
	Beta	70.0	46.0	78.0	82.0	94.0
DEXA scans for younger patients	Alpha	58.6	58.6	44.8	65.5	75.8
	Beta	22.0	22.0	50.0	74.0	86.0
Pediatrics						
Antibiotics pharyngitis	Alpha	42.9	85.7	61.9	90.5	81.0
	Beta	53.6	46.5	53.6	96.5	89.3
Head injury imaging	Alpha	9.5	90.5	85.7	95.3	71.4
	Beta	10.7	35.7	71.5	96.5	89.3
OME referral	Alpha	23.8	52.4	66.6	85.8	90.5
	Beta	21.4	21.4	39.3	78.6	85.7
No cough medications	Alpha	90.5	71.4	47.6	71.5	42.8
	Beta	82.1	60.7	35.7	92.9	78.0
Inhaled steroids for asthma	Alpha	71.4	90.5	81.9	95.2	47.6
	Beta	67.8	78.5	71.5	89.3	85.7

Abbreviations: ECG, electrocardiogram; DEXA, dual energy x-ray absorptiometry; OME, otitis media with effusion; Pap, Papanicolaou.

LIMITATIONS

The items generated by the 3 working groups reflect the opinions of the physicians serving on the working groups. A different group of physicians might have elaborated a different list. We believe that the specific items included on the list may be less important than the process for developing a consensus to change clinical behavior to improve care and reduce risk and cost.

The physicians recruited for field testing may not be representative of physicians in the 3 primary care specialties. A larger sample of physicians selected to reflect the demographic characteristics of the specialties might have responded differently to the survey questions. Resource limitations of this study precluded recruiting a representative sample. Finally, the field testing relied on the opinions of the physicians who participated in the survey rather than on empirical data or actual implementation in practice.

NEXT STEPS

The top-5 lists will be distributed to all NPA physicians in the respective primary care specialties. A virtual practice community will be formed to support physicians' efforts to implement the recommended activities in their practices.

Many of the field testers believed that successful implementation would depend on enlisting patient agreement with the recommendations. Misunderstanding and miscommunication between physicians and patients explain a significant part of why unnecessary and even harmful tests and treatments are ordered. For example, many primary care physicians state that pressure from patients leads them to prescribe antibiotics when they are not indicated. Yet studies have shown that, in fact, patients don't expect antibiotics nearly as often as doctors believe they do.¹⁹⁻²¹

Patient satisfaction and understanding are closely related, and physicians can improve patient satisfaction by focusing on understanding. This can be achieved by acknowledging and validating patient concerns while providing factual information in an easy-to-understand manner, explicitly clarifying the rationale for a selected course of action, and providing a contingency plan that empowers the patient.²²

With this in mind, the NPA is planning to produce training videos to help physicians gain their patients' understanding and support by learning the communication skills necessary to enlist patient partnership in collaborative work. Videos will also be produced specifically for patients, explaining the rationale for the recommendations by clarifying that risks outweigh benefits, and the

Top 5 List in Family Medicine

1. Don't do imaging for low back pain within the first 6 weeks unless red flags* are present

- Imaging of the lumbar spine before 6 weeks does *not* improve outcomes but does increase costs
- Low back pain is the fifth most common reason for all physician visits

* *Red flags include but are not limited to severe or progressive neurological deficits or when serious underlying conditions such as osteomyelitis are suspected*
Sources: AHCPR and Cochrane

2. Don't routinely prescribe antibiotics for acute mild to moderate sinusitis unless symptoms (which must include purulent nasal secretions AND maxillary pain or facial or dental tenderness to percussion) last for 7 or more days OR symptoms worsen after initial clinical improvement

- Most maxillary sinusitis in the ambulatory setting is due to a viral infection that will resolve on its own
- Despite consistent recommendations to the contrary, antibiotics are prescribed in over 80% of outpatient visits for acute sinusitis
- Sinusitis accounts for 16 million office visits and \$5.8 billion in annual healthcare costs

Source: Cochrane and Ann IM

3. Don't order annual ECGs or any other cardiac screening for asymptomatic, low-risk patients

- Little evidence that detection of coronary artery stenosis in asymptomatic patients at low risk for coronary heart disease improves health outcomes
- False-positive tests are likely to lead to harm through unnecessary invasive procedures, over-treatment, and misdiagnosis
- Potential harms of this routine annual screening exceed the potential benefit

Source: USPSTF

4. Don't perform Pap tests on patients younger than 21 years or in women status post hysterectomy for benign disease

- Most dysplasia in adolescents regresses spontaneously; therefore, screening Pap tests done in this age group can lead to unnecessary anxiety, morbidity, and cost
- Pap tests have low yield in women after hysterectomy (for benign disease), and there is poor evidence for improved outcomes

Sources: ACOG (for age), USPSTF (for hysterectomy)

5. Don't use DEXA screening for osteoporosis in women under age 65 years or men under 70 years with no risk factors*

- Not cost-effective in younger, low-risk patients, but cost-effective in older patients

* *Risk factors include but are not limited to fractures after age 50 years, prolonged exposure to corticosteroids, diet deficient in calcium or vitamin D, cigarette smoking, alcoholism, thin and small build*

Sources: NOF, USPSTF, AACE, ACPM

Figure 1. "Top 5" activities in family medicine. AACE indicates American Association of Clinical Endocrinology⁴; ACOG, American College of Obstetrics and Gynecology⁵; ACPM, American College of Preventive Medicine⁶; AHCPR, Agency for Healthcare Policy and Research⁷; Ann IM, *Annals of Internal Medicine*⁸; Cochrane, Cochrane Database of Systematic Reviews⁹; DEXA, dual energy x-ray absorptiometry; ECG, electrocardiogram; NOF, National Osteoporosis Foundation¹⁰; Pap, Papanicolaou; and USPSTF, US Preventive Services Task Force.¹¹

Top 5 List in Internal Medicine

1. Don't do imaging for low back pain within the first 6 weeks unless red flags* are present

- Imaging of the lumbar spine before 6 weeks does *not* improve outcomes but does increase costs
- Low back pain is the fifth most common reason for all physician visits

* *Red flags include but are not limited to severe or progressive neurological deficits or when serious underlying conditions such as osteomyelitis are suspected*
Sources: AHCPR and Cochrane

2. Don't obtain blood chemistry panels (eg, basic metabolic panel) or urinalyses for screening in asymptomatic, healthy adults

- Only lipid screening yielded significant numbers of positive results among asymptomatic patients
- Screen for type 2 diabetes mellitus in asymptomatic adults with hypertension

Source: USPSTF

3. Don't order annual ECGs or any other cardiac screening for asymptomatic, low-risk patients

- Little evidence that detection of coronary artery stenosis in asymptomatic patients at low risk for coronary heart disease improves health outcomes
- False-positive tests are likely to lead to harm through unnecessary invasive procedures, overtreatment, and misdiagnosis
- Potential harms of this routine annual screening exceed the potential benefit

Source: USPSTF

4. Use only generic statins when initiating lipid-lowering drug therapy

- All statins are effective in decreasing mortality, heart attacks, and strokes when does is titrated to effect appropriate LDL cholesterol reduction
- Switch to more expensive brand-name statins (atorvastatin [Lipitor] or rosuvastatin [Crestor]) only if generic statins cause clinical reactions or do not achieve LDL cholesterol goals

Sources: CURVES¹² and MERCURY¹³ trials and metaanalyses

5. Don't use DEXA screening for osteoporosis in women under age 65 years or men under 70 years with no risk factors*

- Not cost-effective in younger, low-risk patients, but cost-effective in older patients

* *Risk factors include but are not limited to fractures after age 50 years, prolonged exposure to corticosteroids, diet deficient in calcium or vitamin D, cigarette smoking, alcoholism, thin and small build*

Sources: NOF, USPSTF, AACE, ACPM

Figure 2. "Top 5" activities in internal medicine. AACE indicates American Association of Clinical Endocrinology⁴; ACOG, American College of Obstetrics and Gynecology⁵; ACPM, American College of Preventive Medicine⁶; AHCPR, Agency for Healthcare Policy and Research⁷; Ann IM, *Annals of Internal Medicine*⁸; Cochrane, Cochrane Database of Systematic Reviews⁹; DEXA, dual energy x-ray absorptiometry; ECG, electrocardiogram; LDL, low-density lipoprotein; NOF, National Osteoporosis Foundation¹⁰; Pap, Papanicolaou; and USPSTF, US Preventive Services Task Force.¹¹ Lipitor is manufactured by Pfizer, New York, New York; Crestor, by AstraZeneca, Wilmington, Delaware.

Top 5 List in Pediatrics

1. Don't prescribe antibiotics for pharyngitis unless the patient tests positive for *streptococcus*

- Most cases of pharyngitis are viral and will not respond to antibiotics, yet antibiotics are prescribed more than half the time
- Antibiotic use has potential risks to the patient, increases bacterial antibiotic resistance, and adds to health care expenses
- The absence of fever, cervical lymphadenopathy, tonsillar exudates, and the presence of cough suggest viral etiology; screening for *streptococcus* may be unnecessary if these criteria are present
- Confirmation of *streptococcus* infection is definitely necessary before antibiotic use can be justified

Sources: AHRQ, Cochrane, EE

2. Don't obtain diagnostic images for minor head injuries without loss of consciousness or other risk factors

- Imaging low-risk patients rarely detects traumatic abnormalities, and of the abnormalities detected, few, if any, require surgery
- Higher risk factors include dizziness, external signs of injury, changes in neurologic function, and dangerous mechanism of injury (eg, bicycle-related injury, fall from 3 feet or more or 5 stairs), age younger than 2 years, Glasgow coma score less than 15, and evidence of basilar skull fracture (eg, "raccoon eyes," hemotympanum)
- Early exposure to radiation poses a significant risk of radiation-attributed cancers – as high as 1 case in 1400 among infants exposed to cranial CT

Source: AAP/AAFP Guidelines

3. Don't refer OME early in the course of the problem

- Many cases of OME resolve spontaneously within 3 months with no adverse consequences
- Reasons for early referral include craniofacial or neurological abnormalities, language delay or learning problems, and when structural abnormalities of the eardrum or middle ear are suspected

Source: AAP/AAFP Guidelines, NICE

4. Advise patients not to use cough and cold medications

- There is little evidence that over-the-counter cough and cold medications reduce cough, rhinorrhea, or shorten the duration of illness. Rather, they can cause adverse consequences including death
- Yet, over 10% of children use a cough and cold medication *every week*

Sources: AAP, Cochrane, FDA

5. Use inhaled corticosteroids to control asthma appropriately

- Use of controlling medication for persistent asthma reduces asthma exacerbations, ED visits, and hospital admissions
- Threshold: More than 4 wheezing episodes *or* 2 episodes requiring oral corticosteroids within 6 months
- Inhaled corticosteroids are relatively safe and well tolerated

Sources: NAEPP

Figure 3. "Top 5" activities in pediatrics. AAP/AAFP indicates the American Academy of Pediatrics/American Academy of Family Practice¹⁴; ACPM, American College of Preventive Medicine⁶; AHCPR, Agency for Healthcare Policy and Research⁷; CT, computed tomography; ED, emergency department; EE, Essential Evidence¹⁵; FDA, US Food and Drug Administration¹⁶; NAEPP, National Asthma Education and Prevention Program¹⁷; NICE, National Institute for Health and Clinical Excellence¹⁸; and OME, otitis media with effusion.

link between overutilization and increases in insurance premiums.

Patient-centered approaches that discuss expectations and share information with patients have been shown to successfully reduce antibiotic prescriptions in primary care.²³ Effective implementation of the top-5 lists must respect patients' values and beliefs while also honoring clinical logic and protecting finite resources.²⁴

The NPA also plans to request the endorsements of consumer groups and patient safety groups for the recommendations in the top-5 lists. Having such endorsements will help dispel the misconception that these clinical recommendations represent rationing and support the idea that often less is truly more.

Accepted for Publication: March 24, 2011.

Published Online: May 23, 2011. doi:10.1001/archinternmed.2011.231

Corresponding Author: Stephen R. Smith, MD, 899 Montauk Ave, New London, CT 06320 (Stephen_R_Smith@brown.edu). Reprints: National Physicians Alliance, 888 16th St NW, Ste 800, Washington, DC 20006.

Author Contributions: All members of The Good Stewardship Working Group are authors and had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

The Good Stewardship Working Group Members: Irene Aguilar, MD, Department of Medicine, University of Colorado Health Sciences Center, Denver; Zackary D. Berger,

MD, Department of Medicine, Johns Hopkins School of Medicine, Baltimore, Maryland; Danielle Casher, MD, Roxborough Pediatrics, Philadelphia, Pennsylvania; Ricky Y. Choi, MD, Department of Pediatrics, University of California, San Francisco; Jonas B. Green, MD, Department of Medicine, University of California, Los Angeles; Elizabeth G. Harding, MD, North County Community Healthcare, Flagstaff, Arizona; Jeffrey R. Jaeger, MD, Division of General Internal Medicine, University of Pennsylvania School of Medicine, Philadelphia; Arthur Lavin, MD, Department of Pediatrics and Advanced Pediatrics, Case Western Reserve University School of Medicine, Cleveland, Ohio; Rebecca Martin, National Physicians Alliance, Washington, DC; Lynda G. Montgomery, MD, Department Family Medicine, Case Western Reserve University School of Medicine; Nancy Morioka-Douglas, MD, Geriatric Education Center, Stanford University School of Medicine, Stanford, California; Judith A. Murphy, MD, retired; Lauren Oshman, MD, Concord, New Hampshire, Family Medicine; Bethany Picker, MD, Family Medicine Residency Program, Central Maine Medical Center, Lewiston; Stephen R. Smith, MD, Department of Family Medicine, Warren Alpert Medical School of Brown University, Providence, Rhode Island; Shubha Venkatesh, MPH, University of Connecticut School of Medicine, Farmington, ; Mozella Williams, MD, Department of Family and Community Medicine, University of Maryland School of Medicine, Baltimore; Gwendolyn M. Wright, MD, Department of Pediatrics, Linda Vista Family Health Center, San Diego, California.

Financial Disclosure: None reported.

Funding/Support: Funding for this project was provided by the American Board of Internal Medicine Foundation.

REFERENCES

1. ABIM Foundation. Medical Professionalism in the New Millenium: A Physician Charter. 2002. <http://www.abimfoundation.org/Professionalism/-/media/Files/Physician%20Charter.aspx>. Accessed April 21, 2011.
2. Brody H. Medicine's ethical responsibility for health care reform—the Top Five list. *N Engl J Med*. 2010;362(4):283-285.
3. Evaluation Research Team, Centers for Disease Control and Prevention. Gaining consensus among stakeholders through the nominal group technique. <http://www.cdc.gov/HealthyYouth/evaluation/pdf/brief7.pdf>. Accessed April 13, 2011.
4. Hodgson SF, Watts NB, Bilezikian JP, et al; AACE Osteoporosis Task Force. American Association of Clinical Endocrinologists medical guidelines for clinical practice for the prevention and treatment of postmenopausal osteoporosis: 2001 edition, with selected updates for 2003. *Endocr Pract*. 2003;9(6):544-564.
5. ACOG Committee on Practice Bulletins—Gynecology. ACOG Practice Bulletin no. 109: Cervical cytology screening. *Obstet Gynecol*. 2009;114(6):1409-1420.
6. Lim LS, Hoeksema LJ, Sherin K; ACPM Prevention Practice Committee. Screening for osteoporosis in the adult U.S. population: ACPM position statement on preventive practice. *Am J Prev Med*. 2009;36(4):366-375.
7. Bigos S, Bowyer O, Braen G, et al. Acute Low Back Problems in Adults: US Dept of Health and Human Services AHCPR Clinical Practice Guideline No. 14: AHCPR Publication No. 95-0642. <http://d4c2.com/d4c2-000038.htm>. Accessed April 21, 2011.
8. Hickner JM, Bartlett JG, Besser RE, Gonzales R, Hoffman JR, Sande MA; American Academy of Family Physicians; American College of Physicians-American Society of Internal Medicine; Centers for Disease Control; Infectious Diseases Society of America. Principles of appropriate antibiotic use for acute rhinosinusitis in adults: background. *Ann Intern Med*. 2001;134(6):498-505.
9. Ahovuo-Saloranta A, Borisenko OV, Kovanen N, et al. Antibiotics for acute maxillary sinusitis. *Cochrane Database Syst Rev*. 2008;2(2):CD000243.
10. National Osteoporosis Foundation. *Clinician's Guide to Prevention and Treatment of Osteoporosis*. Washington, DC: National Osteoporosis Foundation; 2010.
11. U.S. Preventive Services Task Force. Screening for osteoporosis in postmenopausal women: recommendations and rationale. *Ann Intern Med*. 2002;137(6):526-528.
12. Jones P, Kafonek S, Laurora I, Hunninghake D. Comparative dose efficacy study of atorvastatin versus simvastatin, pravastatin, lovastatin, and fluvastatin in patients with hypercholesterolemia (the CURVES study). *Am J Cardiol*. 1998;81(5):582-587.
13. Stender S, Schuster H, Barter P, Watkins C, Kallend D; MERCURY I Study Group. Comparison of rosuvastatin with atorvastatin, simvastatin and pravastatin in achieving cholesterol goals and improving plasma lipids in hypercholesterolaemic patients with or without the metabolic syndrome in the MERCURY I trial. *Diabetes Obes Metab*. 2005;7(4):430-438.
14. Committee on Quality Improvement, American Academy of Pediatrics and Commission on Clinical Policies and Research, American Academy of Family Physicians. The management of minor closed head injury in children. *Pediatrics*. 1999;104(6):1407-1415.
15. Essential Evidence. Sore throat and pharyngitis (child): September 23, 2009 update. <http://www.essentialevidenceplus.com/>. Accessed October 23, 2009.
16. Food and Drug Administration. OTC cough and cold products: not for infants and children under 2 years of age. <http://www.fda.gov/ForConsumers/ConsumerUpdates/ucm048682.htm>. Accessed April 21, 2011.
17. Williams SG, Schmidt DK, Redd SC, Storms W; National Asthma Education and Prevention Program. Key clinical activities for quality asthma care. Recommendations of the National Asthma Education and Prevention Program. *MMWR Recomm Rep*. 2003;52(RR-6):1-8.
18. National Institute for Health and Clinical Excellence. Surgical management of otitis media with effusion in children: NICE clinical guideline 60. <http://www.nice.org.uk/nicemedia/pdf/CG60NICEguideline.pdf>. Accessed April 21, 2011.
19. Britten N, Ukoumunne O. The influence of patients' hopes of receiving a prescription on doctors' perceptions and the decision to prescribe: a questionnaire survey. *BMJ*. 1997;315(7121):1506-1510.
20. Cockburn J, Pit S. Prescribing behaviour in clinical practice: patients' expectations and doctors' perceptions of patients' expectations—a questionnaire study. *BMJ*. 1997;315(7107):520-523.
21. Stivers T, Mangione-Smith R, Elliott MN, McDonald L, Heritage J. Why do physicians think parents expect antibiotics? What parents report vs what physicians believe. *J Fam Pract*. 2003;52(2):140-148.
22. Mangione-Smith R, McGlynn EA, Elliott MN, McDonald L, Franz CE, Kravitz RL. Parent expectations for antibiotics, physician-parent communication, and satisfaction. *Arch Pediatr Adolesc Med*. 2001;155(7):800-806.
23. Altiner A, Brockmann S, Sielk M, Wilm S, Wegscheider K, Abholz HH. Reducing antibiotic prescriptions for acute cough by motivating GPs to change their attitudes to communication and empowering patients: a cluster-randomized intervention study. *J Antimicrob Chemother*. 2007;60(3):638-644.
24. Gwyn R, Elwyn G. When is a shared decision not (quite) a shared decision? Negotiating preferences in a general practice encounter. *Soc Sci Med*. 1999;49(4):437-447.